




LETTER

Meta-population structure and the evolutionary transition to multicellularity

Caroline J. Rose,^{†,¶} 
 Katrin Hammerschmidt,^{‡,¶} 
 Yuriy Pichugin,[§]  and
 Paul B. Rainey* 

Abstract

The evolutionary transition to multicellularity has occurred on numerous occasions, but transitions to complex life forms are rare. Here, using experimental bacterial populations as proxies for nascent multicellular organisms, we manipulate ecological factors shaping the evolution of groups. Groups were propagated under regimes requiring reproduction via a life cycle replete with developmental and dispersal (propagule) phases, but in one treatment lineages never mixed, whereas in a second treatment, cells from different lineages experienced intense competition during the dispersal phase. The latter treatment favoured traits promoting cell growth at the expense of traits underlying group fitness – a finding that is supported by results from a mathematical model. Our results show that the transition to multicellularity benefits from ecological conditions that maintain discreteness not just of the group (soma) phase, but also of the dispersal (germline) phase.

Keywords

Dispersal, fitness decoupling, germ line, group selection, soma.

Ecology Letters (2020) 23: 1380–1390

INTRODUCTION

Multicellular life evolved on independent occasions from single celled ancestral types. Explanations are numerous, ranging from those that emphasise the centrality of cooperation (Queller & Strassmann 2009; Bourke 2011; West *et al.* 2015), to perspectives that give prominence to specific mechanisms (Boraas *et al.* 1998; van Gestel & Tarnita 2017; Herron *et al.* 2019), through those who see vital ingredients residing in ecological factors that underpin emergence of Darwinian properties (Griesemer 2001; Rainey 2007; Godfrey-Smith 2009; Rainey & Kerr 2010; Libby & Rainey 2013a; De Monte & Rainey 2014; Rainey & De Monte 2014; Black *et al.* 2020).

Evidence of a seminal role for ecology comes from an ongoing experiment that took inspiration from ponds studded with reeds and colonised initially with a planktonic-dwelling aerobic microbe. Growth of the microbe depletes oxygen, but the essential resource is available at the air-liquid interface. Growth at the meniscus requires production of adhesive glues (Spiers *et al.* 2002, 2003; Lind *et al.* 2017) that allows formation of mats comprised of sticky cells (simple undifferentiated collectives), but for mats to remain at the surface attachment to a reed is required. Attachment of genetically distinct mats to different reeds ensures variation among mats. From time-to-time a mat detaches from a reed and sinks. Death provides opportunity for an extant mat to export

its success to a fresh reed (provided opportunity for dispersal exists). As a consequence of patchily distributed resources and a means for mats to disperse among reeds, a Darwinian-like process stands to unfold at the level of mats (Rainey & Kerr 2010; Rainey *et al.* 2017; Black *et al.* 2020).

The experimental evolution analogy uses the bacterium *Pseudomonas fluorescens* and glass microcosms as a proxy for reeds (Hammerschmidt *et al.* 2014). Growth of non-sticky (smooth (SM)) planktonic cells depletes oxygen from the broth phase, establishing conditions that favour the evolution of mat-forming (wrinkly spreader (WS)) cells. Formation of mats establishes conditions that favour the further evolution of non-sticky cells within the mat (Rainey & Rainey 2003). Continuing time-lagged frequency dependent interactions between SM and WS types (Rainey & Travisano 1998) generates a simple life cycle (Libby & Rainey 2013b) that becomes the focus of selection (Fig. 1a).

Because the cycle is initially dependent upon spontaneous mutation, it is prone to failure (but lines can also fail through production of fragile mats). Lineages that fail are removed, thus allowing extant types to export their success to new microcosms in precisely the same way as a mat that falls from a reed provides opportunity for competing mats to export their reproductive success. The non-sticky motile cells act as dispersing agents analogous to a germ line. The mat itself serves both an ecological role by ensuring access to oxygen,

New Zealand Institute for Advanced Study, Massey University, Auckland, New Zealand

[†]Present address: Centre d'Ecologie Fonctionnelle et Evolutive, CNRS, Université de Montpellier, Université Paul Valéry Montpellier - IRD - EPHE, 1919 route de Mende, 34293, Montpellier Cedex 5, France

[‡]Present address: Institute of General Microbiology, Kiel University, Kiel, Germany

[§]Present address: Department of Evolutionary Theory, Max Planck Institute for Evolutionary Biology, Plön, Germany

*Correspondence and present address: Department of Microbial Population Biology, Max Planck Institute for Evolutionary Biology, Plön, Germany, and Laboratoire de Génétique de l'Évolution, Chimie Biologie et Innovation, ESPCI Paris, Université PSL, CNRS, Paris, France. E-mail: rainey@evolbio.mpg.de

[¶]These authors contributed equally to this work.

This article has been reviewed & recommended by PCI Evol Biol (<https://doi.org/10.24072pci.evolbiol.100099>)




Figure 1 Experimental Regimes (a) A single ‘mat’ generation consists of a life cycle of two phases. The Maturation Phase is seeded with a single WS cell. SM cells arise within the mat and are harvested after six days of maturation by plating and collection of all SM colonies on agar plates. The SM propagule cells are transferred to a new microcosm to begin a three-day Dispersal Phase, during which WS mat-forming cells arise. At the end of the Dispersal Phase, cells from microcosms are plated once more, and a single WS colony (representative of the most common colony morphology type) is picked to seed the next generation. Mat extinctions occur if there are no SM cells after six days of the Maturation Phase, no WS cells after three days in the Dispersal Phase, or if the mat collapses during the Maturation Phase. (b) and (c) Schematic depiction of a population of eight genetically distinct groups (indicated by different colours) proceeding through one life cycle within their respective non-mixed (b) and mixed (c) ecologies. Figure created with biorender.com.

while also producing seeds for the next generation of mats. In this regard, the mat, in the absence of non-sticky dispersing cells is analogous to soma (and an evolutionary dead-end).

After 10 life cycle generations, mats propagated under the two-phase life cycle regime evolved – in one lineage – a simple genetic switch that reliably transitioned successive life cycle phases, but more striking was the overall impact of the longer timescale (the nine-day time required for doubling of mats) on the shorter timescale (the hourly doubling of cells). As shown by Hammerschmidt *et al.* (2014), selection over the long timescale caused the fitness of mats to increase (as determined by the relative ability of mats to give rise to offspring mats), whereas fitness of the individual cells comprising mats declined (when measured relative to ancestral types). This can be understood in terms of selection over the longer timescale trumping the effects of individual cell selection: over the long-term, successful cells are those whose fitness aligns with the longer timescale defined by the longevity of the nascent multicellular organism (Bourrat 2015; Black *et al.* 2020). Such an alignment of reproductive fates during the transition from cells to multicellular organisms has been referred to as ‘fitness decoupling’ (Michod & Roze 1999) – a term that captures the sense that when selection comes to act over the longer timescale, fitness of the lower level particles ‘decouples’ from that of the higher level collective.

Included in the experiment was a second treatment where mats evolved with a life cycle involving just a single phase: mats gave rise to mat-offspring via a single sticky mat-forming cell. After 10 life cycle generations mat fitness improved, but there was no evidence of fitness decoupling: enhanced fitness of mats was readily explained by enhanced fitness of individual cells (Hammerschmidt *et al.* 2014; Rose *et al.* 2020).

This result drew particular attention to the significance of the two-phase life cycle. For the evolution of multicellular life – given appropriate ecological circumstances – such a life cycle delivers in a single step a second time scale (Black *et al.* 2020) over which selection might act (replete with birth-death events), a developmental programme that stands to become the focus of selection, a reproductive division of labour and even the seeds of a distinction between soma and germ.

One might reasonably ask whether, if such life cycles can arise with such seeming ease, why multicellularity has not arisen more often. One possibility is that ecological conditions are more restrictive than indicated by the reed/pond analogy. In fact, in the regime implemented by Hammerschmidt *et al.* (2014), lineages never mixed: mats were founded by single cells with discreteness maintained by virtue of boundaries afforded by the microcosms, similarly, dispersing cells from each mat were maintained as separate lineages. In the reed/pond analogy, dispersing cells arising from different mats are

released into the planktonic phase and are thus expected to compete with a diverse range of dispersing genotypes. This subtle distinction is likely important.

Here we explore the impact of population structure on the emergence of collective-level individuality. The life cycle from our previously published results ('Non-Mixed Ecology' treatment; Fig. 1b) is contrasted with an identical two-phase life cycle that incorporates competition (mixing) during the dispersal phase. This environmental manipulation, which is here termed the 'Mixed Propagule Ecology' treatment (Fig. 1c), was performed simultaneously with the earlier study. The results show that competition effected during the dispersal phase of a two-stage life cycle leads selection to favour traits that promote cell growth at the expense of traits underlying group fitness. This conflict is due to a trade-off between traits underlying the fitness of groups and their constituent cells, and is supported by findings derived from a mathematical model. While the existence of a germ line can bring about the decoupling of fitness required to achieve a higher level of individuality, intense competition between propagule cells skews selection towards traits that enhance the competitive ability of cells, rather than towards traits that enhance group function, to which the life cycle is integral.

METHODS

Experimental regime

We have previously published the Non-Mixed Ecology treatment in a study that compared its effect relative to a life cycle without reproductive specialisation (Hammerschmidt *et al.* 2014). Here we compare the effect of meta-population structure on the potential for an ETI. Groups of cells ('microcosms') in both the Non-Mixed and Mixed Propagule ecologies of the present study experience identical two-phase life cycles driven by frequency-dependent selection. More specifically, each of the Non-Mixed and Mixed Propagule meta-population ecologies comprised 15 replicates of eight competing groups that were founded with *P. fluorescens* strain SBW25 (Silby *et al.* 2009), and propagated through 10 generations of evolution (one generation equated to one WS-SM-WS life cycle). Further details are described in Fig. 1 and in Appendix S1 in Supporting Information. Fig. 1 contrasts the death–birth process of group competition in the Non-Mixed Ecology, with the physical mixing mode of competition in the Mixed Propagule Ecology.

Fitness assay and life cycle parameters

Cell-level and group-level fitness were assayed after 10 life cycle generations: fifteen representative clones (one per replicate population) were generated from each of the evolved treatments, in addition to 15 ancestral WS lines (each independently isolated from the earliest mats to emerge from the ancestral SM strain SBW25, as described in detail in Hammerschmidt *et al.* 2014). For each genotype, three replicate competition assays were performed in populations of eight microcosms over the timescale of one full life cycle (Fig. 1a)

against a neutrally marked ancestral competitor (Zhang & Rainey 2007). See Appendix S1 for further details.

Our proxy for group-level fitness is the proportion of evolved 'offspring' mats produced at the end of one life cycle by the focal genotype relative to the marked reference strain, and cell-level fitness the total number of cells in the mat at the end of the Maturation Phase. Density of WS and SM cells, and Proportion of SM cells were also assayed at the end of the Maturation Phase. The growth rate of SM cells was determined from three biological replicate SM colonies per line in 96-well microtitre plates shaken at 28°C, and absorbance (OD600) measured in a microplate reader (BioTek) at 24 h. The experiment was repeated three times and the maximum growth rate (V_{max}) was calculated from the maximum slope of absorbance over time. The transition rate between WS and SM cells, that is the level of SM occurrence in the Maturation Phase, and WS occurrence in the Dispersal Phase, was determined in a separate experiment, where static microcosms were individually inoculated with single colonies of the representative WS types. The Maturation Phase was extended from 6 to 12 days, and the Dispersal Phase from 3 to 6 days. At day six of the Maturation Phase, SM cells were collected to inoculate microcosms for the Dispersal Phase. Each day, three replicate microcosms per line were destructively harvested and the presence and number of SM and WS colony forming units recorded.

Statistical analysis

For detecting differences in group-level fitness and transition rate between cells of the evolved and ancestral lines, generalised linear models (error structure: binomial; link function: logit) with the explanatory variables Ecology, and representative clone (nested within Ecology) were calculated. Analyses of variance (ANOVA) were used to test for differences in cell-level fitness, density of WS cells, and density, proportion and growth rate of SM cells between the evolved and ancestral lines. Explanatory variables were Ecology, and representative clone (nested within Ecology). Relationships between the traits and cell and group-level fitness were tested using the mean per representative type accounting for regime. Pearson correlations and regressions were performed.

The sample size was chosen to maximise statistical power and ensure sufficient replication. Assumptions of the tests, that is normality and equal distribution of variances, were visually evaluated. All tests were two-tailed. Effects were considered significant at the level of $P = 0.05$. All statistical analyses were performed with JMP 9. Figures were produced with GraphPad Prism 5.0, Adobe Illustrator CC 17.0.0 and Inkscape 0.92.3.

RESULTS

We begin with a brief description of the contrasting Non-Mixed Ecology and Mixed Propagule Ecology lifecycle regimes. Each generation began with a single WS cell, which through cell-level replication formed a mat at the air-liquid interface (Maturation Phase in Fig. 1a). For a mat to reproduce it was required to be both viable and fecund, that is, it

had to produce SM propagule cells. In both ecological scenarios, competition between groups arose from a death-birth process: following an extinction event, a group was randomly replaced by a surviving competitor group. Extinction/replacement of groups occurred with high frequency [usually due to the lack of SM production (Hammerschmidt *et al.* 2014)], and therefore imposed potent between-group selection.

The two experimental treatments differed solely in manipulation of the Dispersal Phase of the life cycle (Fig. 1b and c). In the Non-Mixed Ecology, SM cells were harvested separately from each surviving group at the end of the Maturation Phase. In contrast, in the Mixed Propagule Ecology, SM cells were harvested and *pooled* from all groups that survived the Maturation Phase. The pooled mixture was then used to seed all eight groups in the Dispersal Phase. In both ecologies, SM propagule cells competed within individual microcosms during the Dispersal Phase to produce WS types, and ultimately for mat formation. At the end of the Dispersal Phase, one colony of the most numerous WS type occurring in each microcosm was transferred to a fresh microcosm to begin the Maturation Phase of the next mat generation. Importantly, this step was performed for both treatments to ensure that all mats at the start of each new generation were seeded from a single cell.

Changes in group and cell fitness

After 10 group generations, changes in both cell and group level fitness were compared with a set of ancestral lines. Given the wide range of mutational pathways for evolution of WS from SM (McDonald *et al.* 2009; Lind *et al.* 2015, 2019), a range of ancestral WS lines was generated for comparison with the evolved lines. Each 'ancestral' line was a WS genotype isolated independently from the first mats emerging from the common SM ancestor (see Methods and (Hammerschmidt *et al.* 2014)) – this enabled a comparison of the distributions of fitness and other parameters of evolved and ancestral lines.

Fitness of all evolved and ancestral lines was estimated by competition in populations of eight microcosms with a common neutrally marked reference SM genotype (Zhang & Rainey 2007) over the timescale of one generation of the mat life cycle (Fig. 1c). The single-celled bottleneck ensured that non-chimeric mat offspring could be counted at the end of the life cycle. Group fitness was the proportion of offspring mats produced (in a population of eight mats) by the focal line relative to the marked competitor, whereas cell fitness was the mean total number of cells present in the microcosms at the end of the Maturation Phase.

Fitness of derived lineages in the Non-Mixed Ecology significantly increased (ability to leave group offspring) relative to the ancestral types ($\chi^2 = 32.660$, d.f. = 1, $P < 0.0001$; Fig. 2a), whereas cell fitness (number of cells present immediately prior to dispersal) decreased ($F_1 = 10.612$, $P = 0.002$; Fig. 2b). In contrast, under the Mixed Propagule Ecology, group fitness did not change ($\chi^2 = 3.137$, d.f.=1, $P = 0.077$; Fig. 2a), whereas cell fitness increased ($F_1 = 56.214$, $P < 0.0001$; Fig. 2b).

At first glance, this is a surprising result. The group (WS mat) phase was identical in both treatments (each group was

founded from a single WS cell). The only difference was the extent of competition among propagule cells. Under the Mixed Propagule Ecology there was no evidence of fitness decoupling as previously reported for the Non-Mixed Ecology (Hammerschmidt *et al.* 2014): there was no change in group fitness (relative to the ancestral type), but fitness of cells increased. Competition among single cells that comprise the propagule phase thus markedly affected the evolutionary fate of the evolving lineages. Such an effect draws attention to the fact that the evolving entities are defined by a life cycle comprised of both soma- and germ-like phases, and not simply by the group (WS) state. In the next sections we unravel the underlying causes, beginning with analysis of the ancestral state.

Trade-off between group and cell fitness

Fig. 2c illustrates a negative relationship between cell and group fitness in the ancestral lines ($\chi^2 = 4.246$, d.f. = 1, $P = 0.0393$). It also shows evidence of a bimodal distribution of group fitness, indicative of a trade-off between traits underpinning cell and group fitness.

Ten generations of selection in the Non-Mixed Ecology shifted the distribution towards the 'high group fitness/ low cell fitness' corner of the graph (Fig. 2d), indicating that group-level selection was more potent than cell-level selection. Under the Mixed Propagule Ecology there was no corresponding change in the relationship between group and cell fitness in the derived lineages (Fig. 2e).

The contrasting responses are most readily understood in terms of differences in the intensity of within- vs. between-lineage selection. In the Non-Mixed Ecology regime lineages do not interact during the Dispersal Phase and thus, competition – wrought via the death and birth of groups – occurred almost exclusively between lineages. Under the Mixed Propagule Ecology regime, whereas the WS mats that initiate the Maturation Phase were discrete and did not mix, the propagules collected after 6 days and used to found the Dispersal Phase were a pooled mixture sampled from each of eight microcosms. Thus, during the Dispersal Phase within-microcosm competition is intense, and appears to have overwhelmed between-lineage competition.

A further factor impacting the Mix-Propagule Ecology, and especially the opportunity for between-lineage selection, was reduced between-lineage variation. This was not directly measured, but was inferred from the identical visual appearance of WS mats in microcosms at end of the Dispersal Phase under the Mixed, but not Non-Mixed, propagule ecologies.

The causes of the reduced between-lineage variation are easily understood and worthy of consideration because they reflect a rarely considered downside of the standard trait group framework (Wilson 1975). Trait group models provide an explanation for the evolution of maintenance of behaviours that are costly to individuals, such as cooperation. Two genotypes are typically assumed: co-operators and defectors. The trait group model assumes that these types are randomly assembled into groups. Within groups, defecting types out-compete co-operators, but groups comprised of co-operators are more productive than groups dominated by defectors.




Figure 2 Changes in group (a) and cell (b) fitness in the Non-Mixed Ecology (Non-Mix) and Mixed Propagule Ecology (Mix) regimes compared to ancestral populations (Anc). Group fitness is the proportion of derived offspring mats after one life cycle relative to a genetically marked reference genotype. Error bars are s.e.m., based on $n = 14$ (Non-Mix) and $n = 15$ (Anc, Mix). ** denotes significance at the level of $P = 0.001$ – 0.01 , and *** at the level of $P < 0.001$. (c–e) Relationship between cell and group fitness in the Non-Mixed (d) and Mixed Propagule (e) Ecologies compared to ancestral (c) populations. Each dot represents the mean of eight lines per replicate population, assessed in three independent competition assays.

Provided there is periodic mixing of the contents of all groups into a single global pool, followed by random assortment into new groups, then cooperation can be maintained. In essence group selection rewards those groups producing the largest numbers of individuals.

In the Mixed Propagule Ecology there is also a significant reward to WS mats that maximise production of SM propagule cells. But it comes with a cost to the efficacy of selection between groups. Consider a single WS type in one of eight microcosms that acquires an early mutation to SM and which therefore yields a vast excess of SM relative to each of the other seven WS mats. This successful SM type is thus over-represented in the pool of SM propagules, which means that each of the eight microcosms that start the Dispersal Phase also contain an excess of this single genotype. Being more numerous, cells of this lineage are likely to be the source of the next WS-causing mutation. Furthermore, mutational biases arising from features of the genotype-to-phenotype map underpinning the transition between SM and WS types (McDonald *et al.* 2009; Lind *et al.* 2015, 2019), means that not only is it likely that the next WS type in each of the eight microcosms arises from the same SM lineage, but also arises via the exact same mutation, or at least a mutation in the

same gene. The overall effect is to eliminate variation between groups, thus essentially eliminating the possibility of between-lineage selection.

Changes in life cycle parameters

To identify traits contributing to differences in fitness between lineages subject to the Non-Mixed and Mixed Propagule ecologies, we measured properties of WS mat and SM propagule cells expected to determine successful multicellular life cycles. After 10 life cycle generations under the Non-Mixed Ecology regime, there was no change in the density, proportion or growth rate of SM cells (measured at the end of the Maturation Phase) (density: $F_1 = 1.278$, $P = 0.2663$; proportion: $F_1 = 2.702$, $P = 0.1095$; growth rate: $F_1 = 2.116$, $P = 0.1522$; Fig. S1a–c), however, the density of WS cells decreased ($F_1 = 8.036$, $P = 0.0065$; Fig. S1d), whereas the rate of transition between WS and SM cells dramatically increased ($\chi^2 = 114.198$, d.f. = 1, $P < 0.0001$; Fig. S1e).

Evolution under the Mixed Propagule Ecology regime led to a reduction in the density and proportion of SM cells (density: $F_1 = 56.214$, $P < 0.0001$; proportion: $F_1 = 102.217$, $P < 0.0001$; growth rate: $F_1 = 2.664$, $P = 0.1103$; Fig. S1a–c),

but an increase in the density of WS cells ($F_1 = 9.904$, $P = 0.0027$; Fig. S1d). Additionally, there was an increase in the rate of transition between WS and SM cells, but this did not approach the magnitude of the effect observed for the Non-Mixed Ecology ($\chi^2 = 12.459$, d.f. = 1, $P = 0.0004$; Fig. S1e).

Understanding the connection between these data and the effects of selection wrought by the two contrasting ecologies is complex. A starting point is to recognise that under both treatment regimens the primary determinant of success is ability of lineages to generate each phase of the life cycle and critically to transition between phases. Given the importance of capacity to transition between states, the dramatic response in the Non-Mixed Ecology is not surprising, however, it is surprising that this response was so reduced in the Mixed Propagule Ecology (Fig. S1e).

As mentioned above, a key difference is the extent of competition between propagules. Under the Mixed Propagule Ecology, propagules arising from mats during the six-day maturation phase must compete directly with propagules derived from other lineages during the dispersal phase. Given that the dispersal phase ends with sampling of a single WS colony (of the most common type) from each microcosm, representation in the next generation is thus determined solely by the number of WS cells at end of the dispersal phase. While this could in principle be achieved by increases in the growth rate or density of SM cells, the selective response was specific to the density of WS cells. In contrast, WS cells arising in the Dispersal Phase of the Non-Mixed Ecology need only outcompete any alternative WS cell types that may (or may not) arise within the same group. Thus, overall, mixing of propagules shifts the emphasis of selection from a developmental programme (capacity to transition through phases of the life cycle), towards density of WS cells (Fig. S1d). Moreover, the fact that under the Mixed Propagule Ecology, transition rate improved only marginally after 10 life cycle generations, whereas WS density significantly increased, points at a trade-off between WS density – and by extension WS growth rate – and ability to transition through phases of the life cycle.

Identification of traits linked to group and cell fitness

The fact that the WS-SM cell transition rate was the only measured parameter to increase in the Non-Mixed Ecology led to recognition that the WS-SM transition rate is associated with group fitness (Fig. 3a–c). Indeed, these two factors are positively correlated in the ancestral lines ($\chi^2 = 28.029$, d.f. = 1, $P < 0.0001$; Fig. 3a). During evolution in the Non-Mixed Ecology, the distribution shifted towards the ‘High Group Fitness/High Transition Rate’ corner of the spectrum with the two parameters still associated ($\chi^2 = 13.657$, d.f. = 1, $P = 0.002$; Fig. 3b).

Cell Fitness in the ancestral lines was strongly associated with the Density of WS cells ($F_1 = 6.673$, $P = 0.023$; Fig. 3d). The distribution of both parameters increased during the Mixed Propagule Ecology ($F_1 = 200.931$, $P < 0.0001$; Fig. 3f) and decreased during the Non-Mixed Ecology ($F_1 = 97.359$, $P < 0.0001$; Fig. 3e).

Trade-off between WS-SM cell transition rate and WS density

A negative relationship (trade-off) exists between WS Density (which is linked to cell fitness) and WS-SM transition rate (which is linked to group fitness) in the ancestral population ($r = -0.705$, $P = 0.003$, $N = 15$; Fig. 3g). The nature of the association explains both the negative relationship between the two levels of fitness observed above (Fig. 2c), and the opposing direction of selection in the two ecologies. While cells were required to survive an identical two-phase life cycle regardless of meta-population structure, these two traits were driven in opposite directions under the two ecologies because of differences in the emphasis of cell and group level selection (Fig. 3h and i).

A simple model embracing cell- and group-level trade-offs

To explore the extent to which the divergent evolutionary trajectories of groups evolving under the non-mixed and mixed regimes might be attributed to the experimentally recognised trade-off between cell and group fitness, and more specifically density of WS cells (the cell-level trait) and transition rate (the group trait), a simple model of group-structured populations was developed (see Extended Methods in Appendix S1 for details).

In the model, cells are characterised by two quantitative traits: growth rate and probability of transitioning between phenotypes. Independent lineages, with parameters drawn randomly from a bivariate normal distribution, found each group. The trade-off between cell and group fitness observed in the ancestral bacterial population (Fig. 2c) was implemented using a trait distribution in which growth rate and transition probability are negatively correlated (lineages comprised of rapidly growing cells tend to transition between phases at a low rate (and vice versa)). During the simulation, lineages passed through the sequence of alternating Maturation and Dispersal Phases separated by sampling bottlenecks. During each phase, lineages grew exponentially until the total cell population of each reached carrying capacity. Additionally, each lineage may switch phenotype, with a probability defined by the corresponding trait value. Lineages that switch are established with the same parameters, but carry the new phenotype. Only lineages that contain a sub-lineage in which the phenotype has switched proceed to the next life cycle phase. Over time, some lineages go extinct due to competition and these are replaced with lineages from the same population. Hence, the distribution of traits across populations changes with time. Evolution was recorded and analysed over 20 full cycles with 600 independent simulations.

The results show that cell growth rate (a proxy for cell fitness) slowly decreased in the Non-Mixed Ecology (Fig. 4a), and rapidly increased in the Mixed Propagule Ecology (Fig. 4b). At the same time, the transition probability (a proxy for group fitness) increased in the Non-Mixed Ecology (Fig. 4c), whereas it remained stable in the Mixed Propagule Ecology (Fig. 4d). Therefore, the model, comprising a minimal model in which evolution affects solely cell growth rate and capacity to switch phenotype, demonstrates that mixed and non-mixed regimes lead to qualitatively different





Figure 3 Relationship between life cycle traits and group and cell fitness. (a–c) Association of transition rate and group fitness in the ancestral populations (a), and in the Non-Mixed (b) and Mixed Propagule (c) Ecologies. (d–f) WS density is positively associated with cell fitness (total number of cells) in the ancestral populations (d), and in the Non-Mixed (e) and Mixed Propagule (f) Ecologies. (g–i) Relationship between WS density and transition rate in the ancestral populations (g), and in the Non-Mixed (h) and Mixed Propagule (i) Ecologies. Group fitness is the proportion of derived offspring mats after one lifecycle relative to a genetically marked reference genotype. Dots represent the mean of eight lines per replicate population, which were assessed in three independent competition assays.

evolutionary outcomes. Additionally, the simulations confirm that the pooling of propagules in the Mixed Propagule Ecology strengthens selection for the trait improving cell fitness (growth rate), which occurs at the expense of traits improving group fitness (transition probability).

Given formulation of the model we asked whether eliminating the trade-off between growth rate and transition

probability affected the response of the evolving lineages to selection. Under the Non-Mixed Ecology, cell growth rate remained essentially unaffected (Fig. S2a), whereas with the trade-off, cell growth rate declined (Fig. 4a). In the Mixed Propagule Ecology, cell growth – in the absence of the trade-off – remained as seen with the trade-off (cf. Fig. 4b with Fig. S2b). Under both non-mixed and mixed regimes

Figure 4 Simulated dynamics of the average cell growth rate in (a) the Non-Mixed Ecology, and (b) the Mixed Propagule Ecology. (c and d) Simulated dynamics of the average transition probability in (c) the Non-Mixed Ecology, and (d) the Mixed Propagule Ecology. Black lines represent median growth/transition rate values across 600 independent realisations of the respective selection regime. Dark grey areas indicate a 50% confidence interval, whereas light grey areas indicate a 95% confidence interval.



transition probability (group fitness) increased although the increase in the mixed regime was less (Fig. S2c and d). Together, results of the simulations are in full agreement with the experimental findings and emphasise the importance of the trade-off between transition rate and WS density as evident in the experimental data.

Summary of principle findings

Table S1 shows differences between ecologies in the partitioning of variation across meta-populations, including downstream consequences, for traits under selection in the non-mixed and mixed regimes. Selection during both phases of the Non-Mixed Ecology favoured a higher WS-SM transition rate. However, under the Mixed Propagule Ecology, the trade-off between WS density and transition rate evident in the ancestral genotype, limited ability of selection to work on the collective lifecycle. Rather than acting on the life cycle as a whole, selection disproportionately affected cell-level selection. Adaptations may arise that allow groups to survive the Maturation Phase of the Mixed Propagule Ecology (*i.e.* high WS-SM transition rate), only to be extinguished during the Dispersal Phase, due to a low competitive ability resulting from reduced WS density. The red box highlights the conflict between the effects of selection on the two incompatible traits involved in the two phases of the life cycle. This is further illustrated in Fig. 5.

DISCUSSION

Life cycles underpin evolutionary transitions to multicellularity (Buss 1987; Rainey 2007; Rainey & Kerr 2010; Hammer-schmidt *et al.* 2014). Life cycles solve the problem of group-level reproduction and shape organismal form (Fig. 6) (Buss 1987; Godfrey-Smith 2009; Rainey & Kerr 2010; Libby & Rainey 2013a; van Gestel & Tarnita 2017). Furthermore, life cycles involving reproductive specialisation provide selection with opportunity to act on something altogether novel – a developmental programme – that likely underpinned the rise of complexity in plants, animals and fungi (Grosberg & Strathmann 2007). Of further and particular significance is that life cycles establish the possibility for selection to operate over timescales longer than that of the doubling time of cells (Black *et al.* 2020). When this is accompanied by a death–birth process over the timescale of the life cycle, then selection over this longer timescale trumps within life cycle selection resulting in the fitness of groups decoupling from fitness of the composite cells. In the long-term, successful groups are composed of cells whose reproductive fate aligns with that of the longer time scale. This is the essence of the evolutionary transition from cells to multicellular life.

It is instructive to place the findings from this study in the context of different modes of group reproduction and consequences for the expected long-term relationship between cell and group fitness. Fig. 6 contrasts reproduction of groups via protected (unmixed) propagule lineages,




Figure 5 Ecological conditions can steer the evolution of traits in opposite directions. A meta-population is depicted by the cloud of cells, where the position of the cloud represents the average fitnesses and the size of the cloud represents the diversity within a meta-population. Group and cell fitness are subjected to a trade-off (dashed line eclipses) and cannot be optimised simultaneously. Arrows indicate the direction of selection applied by Maturation and Dispersal Phases of the life cycle. Both phases selected for increased transition rate in the Non-Mixed Ecology, whereas in the Mixed Propagule Ecology the Dispersal Phase promoted increased cell numbers at the expense of transition rate. The mixing procedure resulted in significantly decreased diversity, which further limited opportunity for adaptive evolution of groups. Figure created with biorender.com.




Figure 6 The origins of life cycles and the notion of fitness decoupling. Mode of group reproduction via (a) fragmentation, (b) a germ line (red) in a highly structured population and (c) a germ line with propagule mixing, affects the emergence of individuality. Mode of group reproduction impacts the relationship between two levels of selection: the cell level (relative to the free-living state), and that of the emerging group. (a) illustrates an example of a group that reproduces by fragmentation where fitness is ‘coupled’: group fitness is a by-product of the fitness of the constituent cells. Larger groups contain more cells and produce more offspring. This holds even when the reproductive life cycle involves a single-celled bottleneck – a feature that is expected to reduce within-group competition. (b) and (c) show examples of groups that reproduce via a life cycle involving two cell types – one soma-like and the other germ-like. Such two-phase life cycles allow possibility for traits determining a necessary developmental programme to evolve independent of the growth rate of cells that comprise the nascent organism. This paves the way for the emergence of new kinds of biological individuals where group fitness ‘decouples’ from cell fitness.

unprotected (mixed) propagule lineages and fragmentation. In the latter, group fitness and cell fitness remain aligned. When propagules never mix, selection at the group level

overwhelms cell-level selection, whereas when propagules mix, selection at the cell-level is the predominate driver of future evolutionary change.

When viewed among the diverse manifestations of multicellular life, non-mixing of propagules appears to be important for groups to begin the evolutionary trajectory towards paradigmatic forms of multicellularity, such as seen in metazoans. When propagules mix, our findings suggest the route towards less integrated forms of multicellularity as seen, for example in the social amoeba, is more likely. Fragmentation of groups by equal division is problematic, resulting in the formation of chimeric organisms rife with cell-level conflict, and tellingly is exceedingly rare among multicellular life.

True slime moulds (Myxomycetes), and social Myxobacteria exhibit sophisticated behaviours such as ‘wolf-pack feeding’ that allow cells to benefit from group-living (Bonner 1998). Cellular slime moulds such as the dictyostelids can form multicellular fruiting bodies when their food supply is exhausted (Strmecki *et al.* 2005). Organisms such as these exhibit rudimentary multicellular life cycles with cellular differentiation, and yet they have remained relatively simple for millions of years. This may be due, at least in part, to ecological factors that maintain a high degree of competition between cells from different groups during the single-cell phases of their respective life cycles. It is also likely that the aggregative mode of group formation (‘coming together’) inhibits the process of selection at the aggregate level, compared to groups that form by growth from propagules (‘staying together’) (Tarnita *et al.* 2013). It is interesting to note that in the experiments presented here, the benefits (to group fitness) of staying together were negated in the Mixed Propagule Ecology, which bear more resemblance to the coming together mode of group organisation during the Dispersal Phase of the life cycle.

If non-mixing among propagules is important for selection to work with potency on groups, then attention turns to environments and ecological circumstances that might ensure discreteness of the reproductive phase. Conceivably certain kinds of structured environments, such as found within soil pores might suffice. An alternate set of possibilities exist in environments where the density of propagules is low. For example in the pond-plus-reed example that inspired our experimental studies, low nutrient levels in the pond may be sufficient to limit between-propagule competition.

Given a period of selection for traits that favour the persistence of groups, more integrated collectives may withstand a less structured ecology. In other words, a structured environment can provide the ecological scaffold necessary to support persistence during an initial period of evolution in which complex adaptations arise and prevail over selection solely for growth rate. Upon removal of the scaffold, features such as boundaries that demarcate groups, would allow collectives to continue to function as evolutionary individuals (Black *et al.* 2020).

Extant multicellular organisms tolerate varying degrees of cell-level selection, as evidenced by the diverse modes of multicellular reproduction that incorporate intense competition at the gamete level. Many plants, for example engage in synchronous seed dispersal – a life cycle not unlike that depicted in Fig. 6c. Cancer is a classic example of lower-level selection occurring in many multicellular organisms that is largely contained by selection at the higher level (cancers

generally arise later in life, after reproduction (Nunney 1999)). In polyandrous animals, sexual selection also occurs at two levels: a higher level with competition between individuals for mating, and a lower level with competition between sperm for fertilisation of eggs within female genital tracts. This lower level has often been shown to account for a large fraction of total variance in male fitness (and hence of the opportunity for selection); for example 46% in red jungle fowl (Collet *et al.* 2012), or 40% in snails (Pélessié *et al.* 2014). Competition between units of the lower level (i.e. germ cells) is extreme in many aquatic invertebrates during broadcast spawning. Here, the animals (higher level) never meet, as sperm and eggs (lower level) are released into the water column, where competition among the gametes for fertilisation takes place.

Given the unknown evolutionary history of organisms that reproduce by life cycles in which there is intense cell-level selection, and the seeming incompatibility of such modes of reproduction with our experimental findings, it is important to recognise that such modes are likely derived and determined by ecological conditions experienced after nascent multicellular forms arose. This draws attention to a possible alternate solution for minimising propagule-level competition that stems from development.

Assuming discreteness of the group phase, and opportunities for group-level dispersal, then collectives that evolve capacity to retain the propagule phase as an integral part of the group, releasing newly created offspring only after the multicellular (albeit immature) state has been achieved, would likely fare well. Such groups would experience minimal between-group selection at the single cell stage and selection would be predominantly group-level. That this mode of reproduction is a feature of paradigmatic forms of multicellularity – along with the fact that germ cells do not typically replicate once produced – likely marks the importance of early developmental innovations for the evolution of complex multicellular life.

ACKNOWLEDGEMENTS

We thank B. Kerr and E. Libby for discussion, P. Bourrat for comments on drafts of the manuscript and the sterling efforts of D. Brisson in his editorial role for *PCI Evol Biol*. The work was directly supported by the Marsden Fund Council from government funding administered by the Royal Society of New Zealand, and in part by grant RFP-12-20.

CONFLICT OF INTEREST DISCLOSURE

The authors of this article declare that they have no financial conflict of interest with the content of this article. PBR is a *PCIEvolBiol* recommender.

AUTHOR CONTRIBUTIONS

K.H., C.J.R and P.B.R. contributed to the conception and design of the study. K.H. and C.J.R. performed research, undertook data analysis and prepared figures. Y.P. designed the numerical model, performed simulations and prepared figures. All authors wrote the paper.

DATA AVAILABILITY STATEMENT

All data are available at <https://zenodo.org/record/3748416#.XpGdTC17Fgg>.

REFERENCES

- Black, A.J., Bourrat, P. & Rainey, P.B. (2020). Ecological scaffolding and the evolution of individuality. *Nat. Ecol. Evol.*, 4, 426–436.
- Bonner, J.T. (1998). The origins of multicellularity. *Integr. Biol.*, 1, 27–36.
- Boraas, M.E., Seale, D.B. & Boxhorn, J.E. (1998). Phagotrophy by a flagellate selects for colonial prey: a possible origin of multicellularity. *Evol. Ecol.*, 12, 153–164.
- Bourke, A.F. (2011). *Principles of Social Evolution*. Oxford University Press, Oxford.
- Bourrat, P. (2015). Levels, time and fitness in evolutionary transitions in individuality. *Phil. Theor. Biol.*, 7, 1–17. <https://doi.org/10.3998/ptb.6959004.0007.001>
- Buss, L.W. (1987). *The Evolution of Individuality*. Princeton: Princeton University Press.
- Collet, J., Richardson, D.S., Worley, K. & Pizzari, T. (2012). Sexual selection and the differential effect of polyandry. *Proc. Natl Acad. Sci. USA*, 109, 8641–8645.
- De Monte, S. & Rainey, P.B. (2014). Nascent multicellular life and the emergence of individuality. *J. Biosci.*, 39, 237–248.
- Godfrey-Smith, P. (2009). *Darwinian Populations and Natural Selection*. Oxford: Oxford University Press.
- Griesemer, J. (2001). The units of evolutionary transition. *Selection*, 1(1–3), 67–80.
- Grosberg, R.K. & Strathmann, R.R. (2007). The evolution of multicellularity: A minor major transition? *Annu. Rev. Ecol. Evol. Syst.*, 38, 621–654.
- Hammerschmidt, K., Rose, C.J., Kerr, B. & Rainey, P.B. (2014). Life cycles, fitness decoupling and the evolution of multicellularity. *Nature*, 515, 75–79.
- Herron, M.D., Borin, J.M., Boswell, J.C., Walker, J., Chen, I.-C.-K., Knox, C.A. et al. (2019). De novo origins of multicellularity in response to predation. *Sci. Rep.*, 9, 2328.
- Libby, E. & Rainey, P.B. (2013a). A conceptual framework for the evolutionary origins of multicellularity. *Phys. Biol.*, 10, 035001.
- Libby, E. & Rainey, P.B. (2013b). Eco-evolutionary feed-back and the tuning of proto-developmental life cycles. *PLoS One*, 8(12), e82274. <https://doi.org/10.1371/journal.pone.0082274>.
- Lind, P.A., Farr, A.D. & Rainey, P.B. (2015). Experimental evolution reveals hidden diversity in evolutionary pathways. *eLife*, 4, 441.
- Lind, P.A., Farr, A.D. & Rainey, P.B. (2017). Convergence in experimental *Pseudomonas* populations. *ISME J.*, 11, 589–600.
- Lind, P.A., Libby, E., Herzog, J. & Rainey, P.B. (2019). Predicting mutational routes to new adaptive phenotypes. *eLife*, 8, 38822.
- McDonald, M.J., Gehrig, S.M., Meintjes, P.L., Zhang, X.X. & Rainey, P.B. (2009). Adaptive divergence in experimental populations of *Pseudomonas fluorescens*. IV. Genetic constraints guide evolutionary trajectories in a parallel adaptive radiation. *Genetics*, 183, 1041–1053.
- Michod, R.E. & Roze, D. (1999). Cooperation and conflict in the evolution of individuality. III. Transitions in the unit of fitness. In: *Mathematical and Computational Biology Computational Morphogenesis, Hierarchical Complexity, and Digital Evolution* (ed. Nehaniv, CL). American Mathematical Society, Providence, RI, pp. 47–92.
- Nunney, L. (1999). Lineage selection and the evolution of multistage carcinogenesis. *Proc. R. Soc. B.*, 266, 493–498.
- Pélissié, B., Jarne, P., Sarda, V. & David, P. (2014). Disentangling precopulatory and postcopulatory sexual selection in polyandrous species. *Evolution*, 68, 1320–1331.
- Queller, D.C. & Strassmann, J.E. (2009). Beyond society: the evolution of organismality. *Phil. Trans. R. Soc. B.*, 364, 3143–3155.
- Rainey, P.B. & De Monte, S. (2014). Resolving conflicts during the evolutionary transition to multicellular life. *Annu. Rev. Ecol. Evol. Syst.*, 45, 599–620.
- Rainey, P.B. & Kerr, B. (2010). Cheats as first propagules: A new hypothesis for the evolution of individuality during the transition from single cells to multicellularity. *BioEssays*, 32, 872–880.
- Rainey, P.B. & Rainey, K. (2003). Evolution of cooperation and conflict in experimental bacterial populations. *Nature*, 425, 72–74.
- Rainey, P.B. & Travisano, M. (1998). Adaptive radiation in a heterogenous environment. *Nature*, 394, 69–72.
- Rainey, P.B., Remigi, P., Farr, A.D. & Lind, P.A. (2017). Darwin was right: where now for experimental evolution? *Curr. Opin. Genet. Dev.*, 47, 102–109.
- Rainey, P.B. (2007). Unity from conflict. *Nature*, 446, 616.
- Rose, C.J., Hammerschmidt, K. & Rainey, P.B. (2020). Experimental evolution of nascent multicellularity: Recognizing a Darwinian transition in individuality. *bioRxiv*. <https://doi.org/10.1101/2020.03.02.973792>
- Silby, M.W., Cerdeño-Tárraga, A.M., Vernikos, G.S., Giddens, S.R., Jackson, R.W., Preston, G.M. et al. (2009). Genomic and genetic analyses of diversity and plant interactions of *Pseudomonas fluorescens*. *Genome Biol.*, 10, R51.
- Spiers, A.J., Kahn, S.G., Bohannon, J., Travisano, M. & Rainey, P.B. (2002). Adaptive divergence in experimental populations of *Pseudomonas fluorescens*. I. Genetic and phenotypic bases of wrinkly spreader fitness. *Genetics*, 161, 33–46.
- Spiers, A.J., Bohannon, J., Gehrig, S.M. & Rainey, P.B. (2003). Biofilm formation at the air-liquid interface by the *Pseudomonas fluorescens* SBW25 wrinkly spreader requires an acetylated form of cellulose. *Mol. Microbiol.*, 50, 15–27.
- Strmecki, L., Greene, D.M. & Pears, C.J. (2005). Developmental decisions in *Dictyostelium discoideum*. *Dev. Biol.*, 284, 25–36.
- Tarnita, C.E., Taubes, C.H. & Nowak, M.A. (2013). Evolutionary construction by staying together and coming together. *J. Theor. Biol.*, 320, 10–22.
- van Gestel, J. & Tarnita, C.E. (2017). On the origin of biological construction, with a focus on multicellularity. *Proc. Natl Acad. Sci. USA*, 114, 11018–11026.
- West, S.A., Fisher, R.M., Gardner, A. & Kiers, E.T. (2015). Major evolutionary transitions in individuality. *Proc. Natl Acad. Sci. USA*, 112, 10112–10119.
- Wilson, D.S. (1975). A theory of group selection. *Proc. Natl Acad. Sci. USA*, 72, 143–146.
- Zhang, X.X. & Rainey, P.B. (2007). Construction and validation of a neutrally-marked strain of *Pseudomonas fluorescens* SBW25. *J. Microbiol. Meth.*, 71, 78–81.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.

Editor, Tim Coulson

Manuscript received 25 May 2020

First decision made 27 May 2020

Manuscript accepted 8 June 2020